

**County:** Lake

**Report Date:** 5/2/2024

### Summary

Conduent HCI's Data Scoring Tool provides a systematic ranking of your county's indicators to help set priorities for community health improvement. To read more about this tool, please see the Data Scoring Tool Methodology document. The Data Scoring Tool Report's summary includes the topics that are considered significant health needs, and indicators for which significant disparities were identified for your county. Following this summary you will find the full Data Scoring Tool results, including all indicator scores, all topic scores, and the indicators within each topic area that are considered a significant health need. Please keep in mind that this report is a snapshot in time, and only reflects the data that are current on Healthy Northeast Ohio as of the date the Data Scoring Tool was run. The findings from the Data Scoring Tool should be used, along with other considerations, such as community input, when prioritizing the health needs of your county.

### Topic Scores

All indicator scores are averaged by topic area, to allow for a higher-level ranking of community needs. Resulting scores range from 0-3, where a higher score indicates a greater level of need. Below are the topic areas with scores of 1.50 or higher in your county. This is the cutoff used for a topic area to be considered a significant health need. Please see the results section below for the indicators that make up each of these topic areas.

| Topic              | Indicators | Score |
|--------------------|------------|-------|
| Other Conditions   | 6          | 1.71  |
| Alcohol & Drug Use | 7          | 1.68  |
| Older Adults       | 28         | 1.51  |
| Cancer             | 18         | 1.50  |

### Identified Disparities

The Data Scoring Tool allows for the identification of significant disparities that may exist by race/ethnicity, age, and gender. Below are the indicators where significant disparities were identified in your county.

| Indicator  | County Distribution |    | Value |    | Target |       |       | Score | Precision | Age   |
|--|---------------------|----|-------|----|--------|-------|-------|-------|-----------|-------|
|  | State               | US | State | US | HP2030 | Local | Trend |       |           |       |
| Workers who Drive Alone to Work                          | 1                   | 2  | 2     | 3  | 1.5    |       | 0     | 1.42  | High      | 55-59 |
| Renters Spending 30% or More of Household Income on Rent | 2                   | 1  | 1     | 0  | 3      |       | 2     | 1.39  | High      | 65+   |
| People 25+ with a Bachelor's Degree or Higher            | 0                   | 0  | 2     | 3  | 1.5    |       | 0     | 0.92  | High      | 65+   |
| People 25+ with a High School Diploma or Higher          | 0                   | 0  | 1     | 1  | 1.5    |       | 0     | 0.42  | High      | 65+   |
| Persons with an Internet Subscription                    | 0                   | 0  | 1     | 1  | 1.5    |       | 0     | 0.42  | High      | 65+   |
| People Living Below Poverty Level                        | 0                   | 0  | 0     | 0  | 1      |       | 1     | 0.33  | High      | 12-17 |

| Indicator   | County Distribution |    | Value |    | Target |       |       | Score | Precision | Gender |
|---|---------------------|----|-------|----|--------|-------|-------|-------|-----------|--------|
|   | State               | US | State | US | HP2030 | Local | Trend |       |           |        |
| Age-Adjusted Death Rate due to Unintentional Injuries     | 2                   | 2  | 2     | 3  | 3      |       | 2     | 2.22  | High      | Male   |
| Age-Adjusted Drug and Opioid-Involved Overdose Death Rate | 2                   | 3  | 1     | 3  | 1.5    |       | 1.5   | 2.00  | Medium    | Male   |
| All Cancer Incidence Rate                                 | 2                   | 2  | 2     | 2  | 1.5    |       | 2     | 1.97  | High      | Male   |
| Oral Cavity and Pharynx Cancer Incidence Rate             | 2                   | 1  | 2     | 2  | 1.5    |       | 2     | 1.81  | High      | Male   |
| Age-Adjusted Death Rate due to Cancer                     | 0                   | 1  | 1     | 2  | 3      |       | 1     | 1.11  | High      | Male   |

| Indicator                                       | County Distribution |    | Value |    | Target |       |       | Score | Precision | Race/Ethnicity   |
|---|---------------------|----|-------|----|--------|-------|-------|-------|-----------|--|
|   | State               | US | State | US | HP2030 | Local | Trend |       |           |  |
| Workers who Drive Alone to Work                 | 1                   | 2  | 2     | 3  | 1.5    |       | 0     | 1.42  | High      | American Indian/Alaska Native  |
| Life Expectancy                                 | 0                   | 1  | 1     | 2  | 1.5    |       | 1.5   | 1.17  | Medium    | Black/African American, non-Hispanic   |
| People 25+ with a Bachelor's Degree or Higher   | 0                   | 0  | 2     | 3  | 1.5    |       | 0     | 0.92  | High      | Black/African American, Other  |
| Premature Death                                 | 0                   | 1  | 0     | 1  | 1.5    |       | 1.5   | 0.83  | Medium    | Black/African American, non-Hispanic   |
| People 65+ Living Below Poverty Level           | 0                   | 0  | 0     | 0  | 1.5    |       | 2     | 0.64  | High      | Black/African American   |
| Per Capita Income                               | 0                   | 0  | 1     | 2  | 1.5    |       | 0     | 0.58  | High      | American Indian/Alaska Native, Black/African American, Hispanic/Latino, Other, Two or More Races |
| People 25+ with a High School Diploma or Higher | 0                   | 0  | 1     | 1  | 1.5    |       | 0     | 0.42  | High      | Black/African American   |
| Persons with an Internet Subscription           | 0                   | 0  | 1     | 1  | 1.5    |       | 0     | 0.42  | High      | Native Hawaiian/Pacific Islander   |
| Children Living Below Poverty Level             | 0                   | 0  | 0     | 0  | 1.5    |       | 1     | 0.36  | High      | Hispanic/Latino, Other, Two or More Races  |
| Families Living Below Poverty Level             | 0                   | 0  | 0     | 0  | 1.5    |       | 1     | 0.36  | High      | Hispanic/Latino  |
| Young Children Living Below Poverty Level       | 0                   | 0  | 0     | 0  | 1.5    |       | 1     | 0.36  | High      | Other  |
| People Living Below Poverty Level               | 0                   | 0  | 0     | 0  | 1      |       | 1     | 0.33  | High      | American Indian/Alaska Native, Black/African American, Hispanic/Latino, Other, Two or More Races |
| Median Household Income                         | 0                   | 0  | 0     | 1  | 1.5    |       | 0     | 0.25  | High      | American Indian/Alaska Native, Black/African American, Other                                     |

### Results

The remaining pages of this report include the full Data Scoring Tool results for your county, including indicator scores, topic scores, and scoring results by topic for topics with scores of 1.50 or higher.

# Healthy Communities Institute Data Scoring Tool



County: Lake  
Cuyahoga (Northeast Ohio)  
Total indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM



| Indicator   | County Distribution |     | Value |     | Target |       | Trend | Score | Precision |
|---|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|
|   | State               | US  | State | US  | HP2030 | Local |       |       |           |
| Alcohol-Impaired Driving Deaths                                 | 3                   | 3   | 3     | 3   | 1.5    |       | 3     | 2.92  | High      |
| Osteoporosis: Medicare Population                               | 3                   | 3   | 3     | 3   | 1.5    |       | 2     | 2.64  | High      |
| People 65+ Living Alone   | 2                   | 3   | 2     | 3   | 1.5    |       | 3     | 2.58  | High      |
| Cervical Cancer Incidence Rate                                  | 1.5                 | 2   | 3     | 3   | 1.5    |       | 3     | 2.50  | Medium    |
| Rheumatoid Arthritis or Osteoarthritis: Medicare Population     | 3                   | 3   | 2     | 3   | 1.5    |       | 2     | 2.47  | High      |
| Age-Adjusted Death Rate due to Falls                            | 3                   | 1.5 | 3     | 1.5 | 1.5    |       | 3     | 2.42  | Medium    |
| Youth not in School or Working                                  | 3                   | 3   | 3     | 3   | 1.5    |       | 1     | 2.36  | High      |
| Median Monthly Owner Costs for Households without a Mortgage    | 3                   | 1.5 | 2     | 2   | 1.5    |       | 3     | 2.33  | Medium    |
| Student-to-Teacher Ratio  | 2                   | 3   | 3     | 3   | 1.5    |       | 1.5   | 2.33  | High      |
| Breast Cancer Incidence Rate                                    | 3                   | 3   | 2     | 2   | 1.5    |       | 2     | 2.31  | High      |
| Cancer: Medicare Population                                     | 3                   | 3   | 2     | 2   | 1.5    |       | 2     | 2.31  | High      |
| Hyperlipidemia: Medicare Population                             | 3                   | 3   | 2     | 2   | 1.5    |       | 2     | 2.31  | High      |
| Primary Care Provider Rate                                      | 2                   | 2   | 3     | 3   | 1.5    |       | 2     | 2.31  | High      |
| Social Associations   | 3                   | 2   | 3     | 2   | 1.5    |       | 2     | 2.31  | High      |
| Age-Adjusted Death Rate due to Unintentional Injuries           | 2                   | 2   | 2     | 3   | 3      |       | 2     | 2.22  | High      |
| Pneumonia Vaccinations: Medicare Population                     | 2                   | 1   | 3     | 3   | 1.5    |       | 2     | 2.14  | High      |
| Adults with Arthritis   | 2                   | 3   | 1.5   | 3   | 1.5    |       | 1.5   | 2.08  | Medium    |
| Adults with Cancer  | 3                   | 2   | 1.5   | 3   | 1.5    |       | 1.5   | 2.08  | Medium    |
| Homeowner Spending-to-Income Ratio                              | 3                   | 3   | 2     | 1.5 | 1.5    |       | 1.5   | 2.08  | Medium    |
| Age-Adjusted Death Rate due to Prostate Cancer                  | 2                   | 2   | 2     | 2   | 3      |       | 2     | 2.06  | High      |
| Age-Adjusted Death Rate due to Suicide                          | 2                   | 1.5 | 3     | 1.5 | 3      |       | 2     | 2.06  | Medium    |
| Workers who Walk to Work  | 2                   | 2   | 3     | 3   | 1.5    |       | 1     | 2.03  | High      |
| Age-Adjusted Death Rate due to Cerebrovascular Disease (Stroke) | 2                   | 1.5 | 1     | 1.5 | 3      |       | 3     | 2.00  | Medium    |
| Age-Adjusted Drug and Opioid-Involved Overdose Death Rate       | 2                   | 3   | 1     | 3   | 1.5    |       | 1.5   | 2.00  | Medium    |
| Median Household Gross Rent                                     | 3                   | 1.5 | 2     | 0   | 1.5    |       | 3     | 2.00  | Medium    |
| Workers Commuting by Public Transportation                      | 0                   | 0   | 3     | 3   | 3      |       | 3     | 2.00  | High      |
| All Cancer Incidence Rate                                       | 2                   | 2   | 2     | 2   | 1.5    |       | 2     | 1.97  | High      |
| COPD: Medicare Population                                       | 1                   | 2   | 2     | 3   | 1.5    |       | 2     | 1.97  | High      |
| Linguistic Isolation  | 3                   | 2   | 3     | 0   | 1.5    |       | 2     | 1.97  | High      |
| Stroke: Medicare Population                                     | 2                   | 2   | 3     | 1   | 1.5    |       | 2     | 1.97  | High      |
| Death Rate due to Drug Poisoning                                | 2                   | 3   | 0     | 3   | 3      |       | 1.5   | 1.92  | High      |
| People 65+ Living Alone (Count)                                 | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 3     | 1.92  | Low       |
| People 65+ Living Below Poverty Level (Count)                   | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 3     | 1.92  | Low       |
| Prostate Cancer Incidence Rate                                  | 2                   | 2   | 1     | 1   | 1.5    |       | 3     | 1.92  | High      |
| Oral Cavity and Pharynx Cancer Incidence Rate                   | 2                   | 1   | 2     | 2   | 1.5    |       | 2     | 1.81  | High      |
| Adults who Binge Drink  | 2                   | 2   | 1.5   | 2   | 1.5    |       | 1.5   | 1.75  | Medium    |
| Annual Ozone Air Quality  | 1.5                 | 3   | 1.5   | 1.5 | 1.5    |       | 1.5   | 1.75  | Low       |
| Mortgaged Owners Median Monthly Household Costs                 | 3                   | 1.5 | 2     | 0   | 1.5    |       | 2     | 1.72  | Medium    |
| Preterm Births  | 2                   | 1.5 | 1     | 1.5 | 3      |       | 2     | 1.72  | Medium    |
| Adults 20+ with Diabetes  | 1                   | 2   | 1.5   | 1.5 | 1.5    |       | 2     | 1.64  | Medium    |
| Atrial Fibrillation: Medicare Population                        | 1                   | 2   | 1     | 2   | 1.5    |       | 2     | 1.64  | High      |
| Depression: Medicare Population                                 | 1                   | 2   | 1     | 2   | 1.5    |       | 2     | 1.64  | High      |
| PBT Released  | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 2     | 1.64  | Low       |
| Recognized Carcinogens Released into Air                        | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 2     | 1.64  | Low       |
| Size of Labor Force   | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 2     | 1.64  | Low       |
| Unemployed Workers in Civilian Labor Force                      | 1                   | 2   | 1     | 2   | 1.5    |       | 2     | 1.64  | High      |
| Weeks of Moderate Drought or Worse                              | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 2     | 1.64  | Low       |
| Gender Pay Gap  | 1.5                 | 2   | 2     | 2   | 1.5    |       | 1     | 1.61  | Medium    |
| College Tuition Spending-to-Income Ratio                        | 2                   | 2   | 1     | 1.5 | 1.5    |       | 1.5   | 1.58  | Medium    |
| Day Care Center and Preschool Spending-to-Income Ratio          | 2                   | 2   | 1     | 1.5 | 1.5    |       | 1.5   | 1.58  | Medium    |
| Home Renter Spending-to-Income Ratio                            | 2                   | 2   | 1     | 1.5 | 1.5    |       | 1.5   | 1.58  | Medium    |
| Insufficient Sleep  | 1                   | 2   | 1     | 2   | 3      |       | 1.5   | 1.58  | High      |
| Non-Physician Primary Care Provider Rate                        | 1                   | 2   | 3     | 3   | 1.5    |       | 0     | 1.58  | High      |
| Preventable Hospital Stays: Medicare Population                 | 2                   | 2   | 2     | 3   | 1.5    |       | 0     | 1.58  | High      |
| Obesity: Medicare Population                                    | 1                   | 1.5 | 0     | 3   | 1.5    |       | 2     | 1.56  | Medium    |
| Ischemic Heart Disease: Medicare Population                     | 1                   | 2   | 2     | 2   | 1.5    |       | 1     | 1.53  | High      |
| Access to Parks   | 0                   | 1.5 | 3     | 1.5 | 1.5    |       | 1.5   | 1.50  | Low       |
| Adults who Smoke  | 0                   | 1   | 1.5   | 3   | 3      |       | 1.5   | 1.50  | Medium    |
| High Blood Pressure Prevalence                                  | 1                   | 1   | 1.5   | 3   | 0      |       | 1.5   | 1.50  | Medium    |
| Mammogram in Past 2 Years: 50-74                                | 1                   | 1   | 1.5   | 2   | 3      |       | 1.5   | 1.50  | Medium    |
| Proximity to Highways   | 3                   | 1.5 | 0     | 1.5 | 1.5    |       | 1.5   | 1.50  | Low       |
| Daily Dose of UV Irradiance                                     | 1                   | 1.5 | 1     | 1.5 | 1.5    |       | 2     | 1.47  | Medium    |
| Age-Adjusted Death Rate due to Coronary Heart Disease           | 1                   | 1.5 | 2     | 1.5 | 3      |       | 1     | 1.44  | Medium    |
| Age-Adjusted Death Rate due to Lung Cancer                      | 1                   | 1   | 1     | 3   | 3      |       | 1     | 1.44  | High      |
| Adult Day Care Spending-to-Income Ratio                         | 2                   | 1   | 1     | 1.5 | 1.5    |       | 1.5   | 1.42  | Medium    |
| Adults who Experienced Coronary Heart Disease                   | 0                   | 1   | 1.5   | 3   | 1.5    |       | 1.5   | 1.42  | Medium    |
| Adults who have had a Routine Checkup                           | 2                   | 1   | 1.5   | 1   | 1.5    |       | 1.5   | 1.42  | Medium    |
| Adults who Have Taken Medications for High Blood Pressure       | 2                   | 1   | 1.5   | 1   | 1.5    |       | 1.5   | 1.42  | Medium    |
| Adults with COPD  | 0                   | 1   | 1.5   | 3   | 1.5    |       | 1.5   | 1.42  | Medium    |
| Adults with Kidney Disease                                      | 1                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.42  | Medium    |
| Age-Adjusted Death Rate due to Diabetes                         | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 3     | 1.42  | Medium    |
| Health Insurance Spending-to-Income Ratio                       | 1                   | 1   | 2     | 1.5 | 1.5    |       | 1.5   | 1.42  | Medium    |

# Healthy Communities Institute Data Scoring Tool



County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM



| Indicator  | County Distribution |     | Value |     | Target |       | Trend | Score | Precision |
|--|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|
|  | State               | US  | State | US  | HP2030 | Local |       |       |           |
| Workers who Drive Alone to Work  | 1                   | 2   | 2     | 3   | 1.5    |       | 0     | 1.42  | High      |
| Adults 20+ Who Are Obese   | 0                   | 2   | 1.5   | 1.5 | 0      |       | 2     | 1.39  | Medium    |
| Grandparents Who Are Responsible for Their Grandchildren                 | 0                   | 1.5 | 0     | 3   | 1.5    |       | 2     | 1.39  | Medium    |
| Renters Spending 30% or More of Household Income on Rent                 | 2                   | 1   | 1     | 0   | 3      |       | 2     | 1.39  | High      |
| 8th Grade Students Proficient in Math                                    | 2                   | 1.5 | 1     | 1.5 | 1.5    |       | 1     | 1.36  | Medium    |
| Number of Extreme Precipitation Days                                     | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 1     | 1.36  | Low       |
| High School Graduation   | 2                   | 1.5 | 1     | 1.5 | 1      |       | 1     | 1.33  | Medium    |
| 4th Grade Students Proficient in English/Language Arts                   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.31  | Medium    |
| Babies with Low Birthweight  | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.31  | Medium    |
| Hypertension: Medicare Population  | 0                   | 1   | 1     | 2   | 1.5    |       | 2     | 1.31  | High      |
| Poor Mental Health: Average Number of Days                               | 0                   | 1   | 1     | 2   | 1.5    |       | 2     | 1.31  | High      |
| Age-Adjusted Death Rate due to Colorectal Cancer                         | 1                   | 1   | 1     | 2   | 3      |       | 1     | 1.28  | High      |
| Tobacco Use: Medicare Population   | 0                   | 1.5 | 1     | 3   | 1.5    |       | 1     | 1.28  | Medium    |
| Adults Ever Diagnosed with Depression                                    | 0                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.25  | Medium    |
| Adults who Experienced a Stroke  | 0                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.25  | Medium    |
| Adults Who Frequently Used Quick Service Restaurants: Past 30 Days       | 2                   | 3   | 1     | 1   | 1.5    |       | 0     | 1.25  | High      |
| Annual Particulate Pollution   | 1.5                 | 0   | 1.5   | 1.5 | 1.5    |       | 1.5   | 1.25  | Low       |
| High Cholesterol Prevalence: Past 5 Years                                | 1                   | 1   | 1.5   | 1   | 1.5    |       | 1.5   | 1.25  | Medium    |
| Poor Mental Health: 14+ Days   | 0                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.25  | Medium    |
| Poor Physical Health: 14+ Days   | 0                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.25  | Medium    |
| Utilities Spending-to-Income Ratio                                       | 1                   | 1   | 1     | 1.5 | 1.5    |       | 1.5   | 1.25  | Medium    |
| Adults who Drink Excessively   | 2                   | 2   | 0     | 1   | 1.5    |       | 1     | 1.19  | High      |
| Alzheimer's Disease or Dementia: Medicare Population                     | 2                   | 1   | 1     | 1   | 1.5    |       | 1     | 1.19  | High      |
| Children in Single-Parent Households                                     | 2                   | 2   | 0     | 1   | 1.5    |       | 1     | 1.19  | High      |
| Gonorrhea Incidence Rate   | 2                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.19  | Medium    |
| Households that are Asset Limited, Income Constrained, Employed (ALICE)  | 1                   | 1.5 | 1     | 1.5 | 1.5    |       | 1     | 1.19  | Medium    |
| Mothers who Received Early Prenatal Care                                 | 1                   | 1.5 | 1     | 1.5 | 1.5    |       | 1     | 1.19  | Medium    |
| Adults who Feel Overwhelmed by Financial Burdens                         | 1                   | 1   | 1     | 1   | 1.5    |       | 1.5   | 1.17  | Medium    |
| Age-Adjusted Death Rate due to Unintentional Poisonings                  | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1.5   | 1.17  | Medium    |
| Life Expectancy  | 0                   | 1   | 1     | 2   | 1.5    |       | 1.5   | 1.17  | Medium    |
| Age-Adjusted Death Rate due to Alzheimer's Disease                       | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.14  | Medium    |
| Age-Adjusted Death Rate due to Kidney Disease                            | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.14  | Medium    |
| Asthma: Medicare Population  | 1                   | 1   | 1     | 0   | 1.5    |       | 2     | 1.14  | High      |
| Dentist Rate   | 0                   | 0   | 1     | 2   | 1.5    |       | 2     | 1.14  | High      |
| Food Environment Index   | 1                   | 1   | 0     | 1   | 1.5    |       | 2     | 1.14  | High      |
| Self-Reported General Health Assessment: Good or Better                  | 0                   | 0   | 1     | 2   | 1.5    |       | 2     | 1.14  | High      |
| Total Employment Change  | 1                   | 2   | 0     | 0   | 1.5    |       | 2     | 1.14  | High      |
| Age-Adjusted Death Rate due to Cancer                                    | 0                   | 1   | 1     | 2   | 3      |       | 1     | 1.11  | High      |
| Children with Health Insurance   | 1.5                 | 1   | 1     | 1   | 1.5    |       | 1     | 1.11  | Medium    |
| Persons without Health Insurance   | 1.5                 | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.11  | Low       |
| Adults who Drank Soft Drinks: Past 7 Days                                | 0                   | 1.5 | 1     | 1   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Adults who Follow a Regular Exercise Routine                             | 0                   | 1.5 | 1     | 1   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Adults with Current Asthma   | 0                   | 1   | 1.5   | 1   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Child Care Centers   | 1                   | 1.5 | 1     | 0   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Cholesterol Test History   | 0                   | 0   | 1.5   | 2   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Cigarette Spending-to-Income Ratio                                       | 0                   | 1   | 1     | 1.5 | 1.5    |       | 1.5   | 1.08  | Medium    |
| Colon Cancer Screening: USPSTF Recommendation                            | 0                   | 1   | 1.5   | 1   | 1.5    |       | 1.5   | 1.08  | Medium    |
| Lung and Bronchus Cancer Incidence Rate                                  | 1                   | 1   | 1     | 3   | 1.5    |       | 0     | 1.08  | High      |
| Student Loan Spending-to-Income Ratio                                    | 1                   | 1   | 0     | 1.5 | 1.5    |       | 1.5   | 1.08  | Medium    |
| Vocational, Technical, and Other School Tuition Spending-to-Income Ratio | 0                   | 1   | 1     | 1.5 | 1.5    |       | 1.5   | 1.08  | Medium    |
| People 65+ Living Below 200% of Poverty Level                            | 1.5                 | 1   | 0     | 0   | 1.5    |       | 2     | 1.06  | Medium    |
| 4th Grade Students Proficient in Math                                    | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| 8th Grade Students Proficient in English/Language Arts                   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| Adults 20+ who are Sedentary   | 0                   | 1   | 1.5   | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| Age-Adjusted Death Rate due to Influenza and Pneumonia                   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| Babies with Very Low Birthweight   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| Liquor Store Density   | 1                   | 0   | 3     | 0   | 1.5    |       | 1     | 1.03  | High      |
| Poor Physical Health: Average Number of Days                             | 0                   | 1   | 1     | 2   | 1.5    |       | 1     | 1.03  | High      |
| Solo Drivers with a Long Commute   | 1                   | 1   | 2     | 0   | 1.5    |       | 1     | 1.03  | High      |
| Violent Crime Rate   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 1.03  | Medium    |
| Adults Who Used Electronic Cigarettes: Past 30 Days                      | 0                   | 1   | 1     | 1   | 1.5    |       | 1.5   | 1.00  | Medium    |
| Households with a Smartphone   | 0                   | 0   | 1     | 2   | 1.5    |       | 1.5   | 1.00  | High      |
| Adults with Health Insurance: 18+  | 0                   | 0   | 1     | 1   | 1.5    |       | 2     | 0.97  | High      |
| Flu Vaccinations: Medicare Population                                    | 0                   | 0   | 1     | 1   | 1.5    |       | 2     | 0.97  | High      |
| Households with Cash Public Assistance Income                            | 1                   | 1   | 0     | 0   | 1.5    |       | 2     | 0.97  | High      |
| Adults with Health Insurance   | 1.5                 | 0   | 1     | 1   | 1.5    |       | 1     | 0.94  | Medium    |
| Mothers who Smoked During Pregnancy                                      | 0                   | 1.5 | 0     | 1.5 | 3      |       | 1     | 0.94  | Medium    |
| Voter Turnout: Presidential Election                                     | 0                   | 1.5 | 1     | 1.5 | 0      |       | 1     | 0.94  | Medium    |
| Adults 65+ who Received Recommended Preventive Services: Males           | 0                   | 0   | 1.5   | 1   | 1.5    |       | 1.5   | 0.92  | Medium    |
| Cervical Cancer Screening: 21-65   | 0                   | 0   | 1.5   | 1   | 1.5    |       | 1.5   | 0.92  | Medium    |
| Gasoline and Other Fuels Spending-to-Income Ratio                        | 0                   | 0   | 1     | 1.5 | 1.5    |       | 1.5   | 0.92  | Medium    |
| Overcrowded Households   | 1                   | 1.5 | 0     | 0   | 1.5    |       | 1.5   | 0.92  | Medium    |

# Healthy Communities Institute Data Scoring Tool



County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM



| Indicator  | County Distribution |     | Value |     | Target |       | Trend | Score | Precision |
|--|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|
|  | State               | US  | State | US  | HP2030 | Local |       |       |           |
| People 25+ with a Bachelor's Degree or Higher                          | 0                   | 0   | 2     | 3   | 1.5    |       | 0     | 0.92  | High      |
| Self-Reported General Health Assessment: Poor or Fair                  | 0                   | 0   | 1.5   | 1   | 1.5    |       | 1.5   | 0.92  | Medium    |
| Families Living Below 200% of Poverty Level                            | 1.5                 | 0   | 0     | 0   | 1.5    |       | 2     | 0.89  | Medium    |
| People Living Below 200% of Poverty Level                              | 1.5                 | 0   | 0     | 0   | 1.5    |       | 2     | 0.89  | Medium    |
| Persons with Health Insurance  | 0                   | 0   | 1     | 1.5 | 1      |       | 1.5   | 0.89  | High      |
| Blood Lead Levels in Children (>=5 micrograms per deciliter)           | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 0.86  | Medium    |
| Diabetes: Medicare Population  | 0                   | 1   | 1     | 1   | 1.5    |       | 1     | 0.86  | High      |
| Heart Failure: Medicare Population                                     | 0                   | 1   | 1     | 1   | 1.5    |       | 1     | 0.86  | High      |
| Households Living Below Poverty Level                                  | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 0.86  | Medium    |
| Households that are Above the Asset Limited, Income Constrained, Emplo | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 0.86  | Medium    |
| Mean Travel Time to Work   | 1                   | 1   | 1     | 0   | 1.5    |       | 1     | 0.86  | High      |
| Students Eligible for the Free Lunch Program                           | 1                   | 0   | 2     | 0   | 1.5    |       | 1     | 0.86  | High      |
| Teen Birth Rate: 15-17   | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 1     | 0.86  | Medium    |
| Access to Exercise Opportunities                                       | 0                   | 0   | 1     | 1   | 1.5    |       | 1.5   | 0.83  | Medium    |
| Adults who Agree Vaccine Benefits Outweigh Possible Risks              | 0                   | 0   | 1     | 1   | 1.5    |       | 1.5   | 0.83  | Medium    |
| Female Population 16+ in Civilian Labor Force                          | 0                   | 0   | 1     | 1   | 1.5    |       | 1.5   | 0.83  | High      |
| Households with a Computer   | 0                   | 0   | 1     | 1   | 1.5    |       | 1.5   | 0.83  | High      |
| Households with a Savings Account                                      | 0                   | 0   | 1     | 1   | 1.5    |       | 1.5   | 0.83  | Medium    |
| Infant Mortality Rate  | 1                   | 1.5 | 0     | 0   | 0      |       | 1.5   | 0.83  | High      |
| Mortgaged Owners Spending 30% or More of Household Income on Hou       | 1.5                 | 0   | 1     | 0   | 0      |       | 1.5   | 0.83  | High      |
| Premature Death  | 0                   | 1   | 0     | 1   | 1.5    |       | 1.5   | 0.83  | Medium    |
| Age-Adjusted Death Rate due to Firearms                                | 1.5                 | 0   | 0     | 0   | 0      |       | 2     | 0.81  | High      |
| Age-Adjusted Death Rate due to Breast Cancer                           | 0                   | 1   | 0     | 1   | 3      |       | 1     | 0.78  | High      |
| Substantiated Child Abuse Rate   | 0                   | 1.5 | 0     | 1.5 | 0      |       | 1     | 0.78  | Medium    |
| Adults 65+ who Received Recommended Preventive Services: Females       | 0                   | 0   | 1.5   | 0   | 1.5    |       | 1.5   | 0.75  | Medium    |
| Adults 65+ with Total Tooth Loss                                       | 0                   | 0   | 1.5   | 0   | 1.5    |       | 1.5   | 0.75  | Medium    |
| Adults without Health Insurance  | 0                   | 0   | 1.5   | 0   | 1.5    |       | 1.5   | 0.75  | Medium    |
| Chlamydia Incidence Rate   | 1                   | 1.5 | 0     | 1.5 | 1.5    |       | 0     | 0.75  | Medium    |
| Home Child Care Spending-to-Income Ratio                               | 0                   | 0   | 0     | 1.5 | 1.5    |       | 1.5   | 0.75  | Medium    |
| Median Household Income: Householders 65+                              | 0                   | 1   | 1     | 2   | 1.5    |       | 0     | 0.75  | High      |
| Mental Health Provider Rate  | 0                   | 0   | 2     | 2   | 1.5    |       | 0     | 0.75  | High      |
| Adults who Visited a Dentist   | 0                   | 0   | 1     | 1   | 1.5    |       | 1     | 0.69  | High      |
| Adults with Internet Access  | 0                   | 0   | 1     | 1   | 1.5    |       | 1     | 0.69  | High      |
| Child Food Insecurity Rate   | 1                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.69  | High      |
| Food Insecurity Rate   | 0                   | 1   | 0     | 1   | 1.5    |       | 1     | 0.69  | High      |
| Mammography Screening: Medicare Population                             | 0                   | 0   | 1     | 1   | 1.5    |       | 1     | 0.69  | High      |
| Population 16+ in Civilian Labor Force                                 | 0                   | 0   | 1     | 1   | 1.5    |       | 1     | 0.69  | High      |
| Adults with Disability Living in Poverty                               | 0                   | 1   | 0     | 0   | 1.5    |       | 1.5   | 0.67  | High      |
| Income Inequality  | 1                   | 0   | 0     | 0   | 1.5    |       | 1.5   | 0.67  | High      |
| Tuberculosis Incidence Rate  | 0                   | 1.5 | 0     | 0   | 0      |       | 1.5   | 0.67  | High      |
| People 65+ Living Below Poverty Level                                  | 0                   | 0   | 0     | 0   | 1.5    |       | 2     | 0.64  | High      |
| Children Living Below 200% of Poverty Level                            | 1.5                 | 0   | 0     | 0   | 1.5    |       | 1     | 0.61  | Medium    |
| Age-Adjusted Death Rate due to Chronic Lower Respiratory Diseases      | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 0     | 0.58  | Medium    |
| Age-Adjusted Death Rate due to Heart Attack                            | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 0     | 0.58  | Medium    |
| Age-Adjusted Death Rate due to Motor Vehicle Collisions                | 1.5                 | 1.5 | 0     | 0   | 1.5    |       | 0     | 0.58  | Medium    |
| Per Capita Income  | 0                   | 0   | 1     | 2   | 1.5    |       | 0     | 0.58  | High      |
| Adults Who Used Smokeless Tobacco: Past 30 Days                        | 0                   | 0   | 0     | 1   | 1.5    |       | 1     | 0.53  | High      |
| Chronic Kidney Disease: Medicare Population                            | 0                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.53  | High      |
| Households without a Vehicle   | 0                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.53  | High      |
| Houses Built Prior to 1950   | 0                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.53  | High      |
| Households with an Internet Subscription                               | 0                   | 0   | 1     | 1   | 1.5    |       | 0     | 0.42  | High      |
| Households with One or More Types of Computing Devices                 | 0                   | 0   | 1     | 1   | 1.5    |       | 0     | 0.42  | High      |
| People 25+ with a High School Diploma or Higher                        | 0                   | 0   | 1     | 1   | 1.5    |       | 0     | 0.42  | High      |
| Persons with an Internet Subscription                                  | 0                   | 0   | 1     | 1   | 1.5    |       | 0     | 0.42  | High      |
| Residential Segregation - Black/White                                  | 0                   | 2   | 0     | 0   | 1.5    |       | 0     | 0.42  | High      |
| Children Living Below Poverty Level                                    | 0                   | 0   | 0     | 0   | 1.5    |       | 1     | 0.36  | High      |
| Families Living Below Poverty Level                                    | 0                   | 0   | 0     | 0   | 1.5    |       | 1     | 0.36  | High      |
| Homeownership  | 0                   | 0   | 0     | 0   | 1.5    |       | 1     | 0.36  | High      |
| Young Children Living Below Poverty Level                              | 0                   | 0   | 0     | 0   | 1.5    |       | 1     | 0.36  | High      |
| People Living Below Poverty Level                                      | 0                   | 0   | 0     | 0   | 1      |       | 1     | 0.33  | High      |
| Colorectal Cancer Incidence Rate                                       | 0                   | 0   | 0     | 1   | 1.5    |       | 0     | 0.25  | High      |
| Median Household Income  | 0                   | 0   | 0     | 1   | 1.5    |       | 0     | 0.25  | High      |
| Severe Housing Problems  | 1                   | 0   | 0     | 0   | 1.5    |       | 0     | 0.25  | High      |
| Homeowner Vacancy Rate   | 0                   | 0   | 0     | 0   | 1.5    |       | 0     | 0.08  | High      |

# Healthy Communities Institute Data Scoring Tool



County: Lake  
Cuyahoga (Northeast Ohio)  
Total indicators: 208  
Thursday 2nd of May 2024 06:03:09 AM

| Topic                               | Indicators | Score |
|-------------------------------------|------------|-------|
| Other Conditions                    | 6          | 1.71  |
| Alcohol & Drug Use                  | 7          | 1.68  |
| Older Adults                        | 28         | 1.51  |
| Cancer                              | 18         | 1.50  |
| Women's Health                      | 6          | 1.45  |
| Heart Disease & Stroke              | 15         | 1.44  |
| Mortality Data                      | 25         | 1.40  |
| Prevention & Safety                 | 7          | 1.34  |
| Diabetes                            | 3          | 1.31  |
| Physical Activity                   | 6          | 1.31  |
| Mental Health & Mental Disorders    | 9          | 1.30  |
| Immunizations & Infectious Diseases | 9          | 1.22  |
| Education                           | 14         | 1.21  |
| Health Care Access & Quality        | 14         | 1.19  |
| Environmental Health                | 20         | 1.18  |
| Wellness & Lifestyle                | 10         | 1.18  |
| Nutrition & Healthy Eating          | 3          | 1.16  |
| Respiratory Diseases                | 13         | 1.15  |
| Community                           | 44         | 1.14  |
| Maternal, Fetal & Infant Health     | 7          | 1.13  |
| Oral Health                         | 4          | 1.10  |
| Tobacco Use                         | 6          | 1.08  |
| Economy                             | 53         | 1.03  |
| Children's Health                   | 6          | 0.88  |
| Health Information Technology       | 6          | 0.63  |

Healthy Communities Institute  
Data Scoring Tool

Scoring Results by Topic

County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM

Topic Selection: **Other Conditions** \*\*\*\*

Other Conditions in Lake County

| Indicator   | County Distribution |     | Value |     | Target |       | Trend | Score | Precision | Disparity Exists |
|---|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|------------------|
|   | State               | US  | State | US  | HP2030 | Local |       |       |           |                  |
| Osteoporosis: Medicare Population                           | 3                   | 3   | 3     | 3   | 1.5    |       | 2     | 2.64  | High      | N/A              |
| Rheumatoid Arthritis or Osteoarthritis: Medicare Population | 3                   | 3   | 2     | 3   | 1.5    |       | 2     | 2.47  | High      | N/A              |
| Adults with Arthritis                                       | 2                   | 3   | 1.5   | 3   | 1.5    |       | 1.5   | 2.08  | Medium    | N/A              |
| Adults with Kidney Disease                                  | 1                   | 1   | 1.5   | 2   | 1.5    |       | 1.5   | 1.42  | Medium    | N/A              |
| Age-Adjusted Death Rate due to Kidney Disease               | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.14  | Medium    | N/A              |
| Chronic Kidney Disease: Medicare Population                 | 0                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.53  | High      | N/A              |

Healthy Communities Institute  
Data Scoring Tool

Scoring Results by Topic

County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM

Topic Selection: Alcohol & Drug Use \*\*\*\*\*

Alcohol & Drug Use in Lake County

| Indicator   | County Distribution |     | Value |     | Target |       | Trend | Score | Precision | Disparity Exists |
|---|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|------------------|
|   | State               | US  | State | US  | HP2030 | Local |       |       |           |                  |
| Alcohol-impaired Driving Deaths                           | 3                   | 3   | 3     | 3   | 1.5    |       | 3     | 2.92  | High      | N/A              |
| Age-Adjusted Drug and Opioid-Involved Overdose Death Rate | 2                   | 3   | 1     | 3   | 1.5    |       | 1.5   | 2     | Medium    | Y                |
| Death Rate due to Drug Poisoning                          | 2                   | 3   | 0     | 3   | 3      |       | 1.5   | 1.92  | High      | N                |
| Adults who Binge Drink                                    | 2                   | 2   | 1.5   | 2   | 1.5    |       | 1.5   | 1.75  | Medium    | N/A              |
| Adults who Drink Excessively                              | 2                   | 2   | 0     | 1   | 1.5    |       | 1     | 1.19  | High      | N/A              |
| Liquor Store Density                                      | 1                   | 0   | 3     | 0   | 1.5    |       | 1     | 1.03  | High      | N/A              |
| Mothers who Smoked During Pregnancy                       | 0                   | 1.5 | 0     | 1.5 | 3      |       | 1     | 0.94  | Medium    | N/A              |

Healthy Communities Institute  
Data Scoring Tool

Scoring Results by Topic

County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM

Topic Selection: Older Adults  
Topic Score: 3.51 \*\*\*\*\*

Older Adults in Lake County

| Indicator  | County Distribution |     | Value |     | Target |       | Trend | Score | Precision | Disparity Exists |
|--|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|------------------|
|  | State               | US  | State | US  | HP2030 | Local |       |       |           |                  |
| Osteoporosis: Medicare Population                                | 3                   | 3   | 3     | 3   | 1.5    |       | 2     | 2.64  | High      | N/A              |
| People 65+ Living Alone  | 2                   | 3   | 2     | 3   | 1.5    |       | 3     | 2.58  | High      | N/A              |
| Rheumatoid Arthritis or Osteoarthritis: Medicare Population      | 3                   | 3   | 2     | 3   | 1.5    |       | 2     | 2.47  | High      | N/A              |
| Age-Adjusted Death Rate due to Falls                             | 3                   | 1.5 | 3     | 1.5 | 1.5    |       | 3     | 2.42  | Medium    | N/A              |
| Cancer: Medicare Population                                      | 3                   | 3   | 2     | 2   | 1.5    |       | 2     | 2.31  | High      | N/A              |
| Hyperlipidemia: Medicare Population                              | 3                   | 3   | 2     | 2   | 1.5    |       | 2     | 2.31  | High      | N/A              |
| COPD: Medicare Population  | 1                   | 2   | 2     | 3   | 1.5    |       | 2     | 1.97  | High      | N/A              |
| Stroke: Medicare Population                                      | 2                   | 2   | 3     | 1   | 1.5    |       | 2     | 1.97  | High      | N/A              |
| People 65+ Living Alone (Count)                                  | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 3     | 1.92  | Low       | N/A              |
| People 65+ Living Below Poverty Level (Count)                    | 1.5                 | 1.5 | 1.5   | 1.5 | 1.5    |       | 3     | 1.92  | Low       | N                |
| Prostate Cancer Incidence Rate                                   | 2                   | 2   | 1     | 1   | 1.5    |       | 3     | 1.92  | High      | N                |
| Atrial Fibrillation: Medicare Population                         | 1                   | 2   | 1     | 2   | 1.5    |       | 2     | 1.64  | High      | N/A              |
| Depression: Medicare Population                                  | 1                   | 2   | 1     | 2   | 1.5    |       | 2     | 1.64  | High      | N/A              |
| Ischemic Heart Disease: Medicare Population                      | 1                   | 2   | 2     | 2   | 1.5    |       | 1     | 1.53  | High      | N/A              |
| Adult Day Care Spending-to-Income Ratio                          | 2                   | 1   | 1     | 1.5 | 1.5    |       | 1.5   | 1.42  | Medium    | N/A              |
| Hypertension: Medicare Population                                | 0                   | 1   | 1     | 2   | 1.5    |       | 2     | 1.31  | High      | N/A              |
| Alzheimer's Disease or Dementia: Medicare Population             | 2                   | 1   | 1     | 1   | 1.5    |       | 1     | 1.19  | High      | N/A              |
| Age-Adjusted Death Rate due to Alzheimer's Disease               | 0                   | 1.5 | 0     | 1.5 | 1.5    |       | 2     | 1.14  | Medium    | N/A              |
| Asthma: Medicare Population                                      | 1                   | 1   | 1     | 0   | 1.5    |       | 2     | 1.14  | High      | N/A              |
| Adults 65+ who Received Recommended Preventive Services: Males   | 0                   | 0   | 1.5   | 1   | 1.5    |       | 1.5   | 0.92  | Medium    | N/A              |
| Diabetes: Medicare Population                                    | 0                   | 1   | 1     | 1   | 1.5    |       | 1     | 0.86  | High      | N/A              |
| Heart Failure: Medicare Population                               | 0                   | 1   | 1     | 1   | 1.5    |       | 1     | 0.86  | High      | N/A              |
| Adults 65+ who Received Recommended Preventive Services: Females | 0                   | 0   | 1.5   | 0   | 1.5    |       | 1.5   | 0.75  | Medium    | N/A              |
| Adults 65+ with Total Tooth Loss                                 | 0                   | 0   | 1.5   | 0   | 1.5    |       | 1.5   | 0.75  | Medium    | N/A              |
| Median Household Income: Householders 65+                        | 0                   | 1   | 1     | 2   | 1.5    |       | 0     | 0.75  | High      | N/A              |
| Mammography Screening: Medicare Population                       | 0                   | 0   | 1     | 1   | 1.5    |       | 1     | 0.69  | High      | N/A              |
| People 65+ Living Below Poverty Level                            | 0                   | 0   | 0     | 0   | 1.5    |       | 2     | 0.64  | High      | Y                |
| Chronic Kidney Disease: Medicare Population                      | 0                   | 1   | 0     | 0   | 1.5    |       | 1     | 0.53  | High      | N/A              |



Healthy Communities Institute  
Data Scoring Tool

Scoring Results by Topic

County: Lake  
Cuyahoga (Northeast Ohio)  
Total Indicators: 208  
Thursday 2nd of May 2024 06:02:13 AM

Topic Selection: Cancer  
Topic Score: 1.5 \*\*\*\*\*

Cancer in Lake County

| Indicator  | County Distribution |    | Value |    | Target |       | Trend | Score | Precision | Disparity Exists |
|--|---------------------|----|-------|----|--------|-------|-------|-------|-----------|------------------|
|  | State               | US | State | US | HP2030 | Local |       |       |           |                  |
| Cervical Cancer Incidence Rate                   | 1.5                 | 2  | 3     | 3  | 1.5    |       | 3     | 2.5   | Medium    | N                |
| Breast Cancer Incidence Rate                     | 3                   | 3  | 2     | 2  | 1.5    |       | 2     | 2.31  | High      | N                |
| Cancer: Medicare Population                      | 3                   | 3  | 2     | 2  | 1.5    |       | 2     | 2.31  | High      | N/A              |
| Adults with Cancer                               | 3                   | 2  | 1.5   | 3  | 1.5    |       | 1.5   | 2.08  | Medium    | N/A              |
| Age-Adjusted Death Rate due to Prostate Cancer   | 2                   | 2  | 2     | 2  | 3      |       | 2     | 2.06  | High      | N                |
| All Cancer Incidence Rate                        | 2                   | 2  | 2     | 2  | 1.5    |       | 2     | 1.97  | High      | Y                |
| Prostate Cancer Incidence Rate                   | 2                   | 2  | 1     | 1  | 1.5    |       | 3     | 1.92  | High      | N                |
| Oral Cavity and Pharynx Cancer Incidence Rate    | 2                   | 1  | 2     | 2  | 1.5    |       | 2     | 1.81  | High      | Y                |
| Mammogram in Past 2 Years: 50-74                 | 1                   | 1  | 1.5   | 2  | 3      |       | 1.5   | 1.5   | Medium    | N/A              |
| Age-Adjusted Death Rate due to Lung Cancer       | 1                   | 1  | 1     | 3  | 3      |       | 1     | 1.44  | High      | N                |
| Age-Adjusted Death Rate due to Colorectal Cancer | 1                   | 1  | 1     | 2  | 3      |       | 1     | 1.28  | High      | N                |
| Age-Adjusted Death Rate due to Cancer            | 0                   | 1  | 1     | 2  | 3      |       | 1     | 1.11  | High      | Y                |
| Colon Cancer Screening: USPSTF Recommendation    | 0                   | 1  | 1.5   | 1  | 1.5    |       | 1.5   | 1.08  | Medium    | N/A              |
| Lung and Bronchus Cancer Incidence Rate          | 1                   | 1  | 1     | 3  | 1.5    |       | 0     | 1.08  | High      | N                |
| Cervical Cancer Screening: 21-65                 | 0                   | 0  | 1.5   | 1  | 1.5    |       | 1.5   | 0.92  | Medium    | N/A              |
| Age-Adjusted Death Rate due to Breast Cancer     | 0                   | 1  | 0     | 1  | 3      |       | 1     | 0.78  | High      | N                |
| Mammography Screening: Medicare Population       | 0                   | 0  | 1     | 1  | 1.5    |       | 1     | 0.69  | High      | N/A              |
| Colorectal Cancer Incidence Rate                 | 0                   | 0  | 0     | 1  | 1.5    |       | 0     | 0.25  | High      | N                |